REFERENCES

- Agha-Hosseini M, Kashani L, Aleyaseen A, Ghoreishi A, Rahmanpour H, Zarrinara AR, Akhondzadeh S, (2008), Crocus sativus L. (saffron) in the treatment of premenstrual syndrome: a double-blind, randomised and placebo-controlled trial, BJOG.115., 515-9.
- Akash A., Savaliya, R. P., Shah, B.P., Saranjit Singh, (2010), Screening of Indian aphrodisiac ayurvedic/herbal healthcare products for adulteration with sildenafil, tadalafil and/or vardenafil using LC/PDA and extracted ion LC–MS/TOF, Journal of Pharmaceutical and Biomedical Analysis, 52., 406–409.
- Akhondzadeh S, Tahmacebi-Pour N, Noorbala AA, Amini H, Fallah-Pour H, Jamshidi AH, Khani M. (2005). Crocus sativus L. in the treatment of mild to moderate depression: a double-blind, randomized and placebo-controlled trial. Phytother Res. 19., 148-51.
- Singh A, Kala S, Kapoor D. N., Gupta R., Antar Virk, Samarjeet Singh and Jyoti Chaudhary, (2011), Effect on human sperm mitochondrial activity by Piper betle and Calendula officinalis, Annals of Biological Research, 2., 622-627
- Bastiaan J. Venhuis, Leonore Blok-Tip, Dries de Kaste, (2008), Designer drugs in herbal aphrodisiacs, Forensic Science International, 177., 25–27.
- Busse W. (2000), the significance of quality for efficacy and safety of herbal medicinal products. Drug Information Journal, 34., 15-23.
- C.A. Calliste, D. Kozlowski, J.L. Duroux, Y. Champavier, A.J. Chulia, P. Trouillas, (2010) A new antioxidant from wild nutmeg, Food Chemistry, 118., 489–496.
- Derby CA, Mohr BA, Goldstein I, Feldman HA, Johannes CB, McKinlay JB. (2000), Modifiable risk factors and erectile dysfunction: can lifestyle changes modify risk? Urology, 56., 302–6.
- Gauthaman K., Adaikan P. G. and Prasad R. N. V., (2002), Aphrodisiac properties of *Tribulus Terrestris* extract (Protodioscin) in normal and castrated rats, Life sciences, 71., 1385-1396.
- Esposito K, Giugliano F, Di Palo C. (2004), Effect of lifestyle changes on erectile dysfunction in obese men: a randomized controlled trial. JAMA, 291., 2978–84.

- Jackson G., Boon N., Eardley I., Kirby M., (2010), Erectile dysfunction and coronary artery disease prediction: evidence-based guidance and consensus, Int J Clin Pract, 64., 831–2.
- Hatzimouratidis K, Burnett AL, Hatzichristou D, McCullough AR, Montorsi F, Mulhall JP. (2009), Phosphodiesterase type 5 inhibitors in post prostatectomy erectile dysfunction: a critical analysis of the basic science rationale and clinical application. Eur Urol, 55., 334–47.
- Guohuaa Hu, Yanhuab Lu, Renganga Mao, Dongzhib Wei, Zhengzhia Ma, Zhang Huaa, (2009) , Aphrodisiac properties of Allium tuberosum seeds extract, Journal of Ethnopharmacology, 122 ., 579–582.
- Gauthaman K., Adaikan P.G., Prasad R.N.V., (2002), Aphrodisiac properties of Tribulus Terrestris extract (Protodioscin) in normal and castrated rats, Life Sciences, 71., 1385–1396.
- Kamalipour M, Jamshidi A.H, Akhondzadeh S, (2010), Antidepressant Effect of Crocus sativus: an Evidence Based Review, Winter Journal of Medicinal Plants , 9., 1-4.
- Kanjwani DG, Marathe TP, Chiplunkar SV, Sathaye SSS, (2008), Evaluation of immunomodulatory activity of methanolic extract of Piper betel, J Immunol, 67., 589-93.
- Misra L. and Wagner H., (2004), Alkaloidal constituents of Mucuna pruriens seeds, Phytochemistry, 65., 2565–2567.
- Carro-Jua´reza M., Cervantesa E., Cervantes-Me´ndeza M., Rodrı´guez-Manzob G., (2004), Aphrodisiac properties of Montanoa tomentosa aqueous crude extract in male rats Pharmacology, Biochemistry and Behavior, 78., 129–134.
- Malviya N, Jain S, Bihari V, Vyas S. (2011), Recent studies on aphrodisiac herbs for the management of male sexual dysfunction a review. Acta Poloniae Pharmaceutica Drug Research, 68., 3-8.
- Thakur M., Chauhan N, Bhargava V, Dixit A. (2009), A Comparative Study on Aphrodisiac Activity of Some Ayurvedic Herbs in Male Albino Rats, rich Sex Behav, 38., 1009–1015.
- Miguel C, Isaac L, Octavio B, Alejandra E, (2006), Pro-ejaculatory effect of the aqueous crude extract of cihuapatli (Montanoa tomentosa) in spinal male rats, Journal of Ethnopharmacology, 106., 111–116.

- Choudhary M, Fatima N, Abbasi M, Jalil S, Uddin A, Atta-ur-Rahman, (2004), Phenolic glycosides, a new class of human recombinant nucleotide pyrophosphatase phosphodiesterase-1 inhibitors, Bioorganic & Medicinal Chemistry, 12., 5793–5798.
- Musa T, Musbau A, Adenike T, Ayoade A. (2008), Androgenic potentials of aqueous extract of Massularia acuminata (G. Don) Bullock ex Hoyl. stem in male Wistar rats, Journal of Ethnopharmacology ,118., 508–513
- Chauhan NS, Rao CV, Dixit VK, (2007), Effect of Curculigo orchioides rhizomes on sexual behaviour of male rats, Fitoterapia ,78., 530–534.
- Malviya N, Jain S, Bihari V and Vyas S, (2011), recent studies on aphrodisiac herbs for the management of male sexual dysfunction and a review, Acta Poloniae Pharmaceutica Drug Research, 68., 3-8
- Hung P, Thi V, Hu Won K, Jooyoung C, Sang K, Kwang-iI K, Dae Bang S, Sang J, Won K, (2010), AMP-activated protein kinase (AMPK) activators from Myristica fragrans (nutmeg) and their anti-obesity effect, Bioorganic & Medicinal Chemistry Letters, 20., 4128–4131.
- Prapapan T, Jarupa V, Nanteetip L, Chawlada S, Anchana K, Khanit S, Kornkanok I, (2008), Screening for phosphodiesterase inhibitory activity of Thai medicinal plants, Journal of Ethnopharmacology, 119., 214–217.
- Ramachandran S, Sridhar Y, Kishore S, Saravanan M. Leonard T, Anbalagan N and Sridhar SK, (2004), Aphrodisiac activity of Butea frondosa Koen. ex Roxb. extract in male rats, Phytomedicine, 11., 165–168.
- Sapna S and Ravi TK. (2007), Approaches towards development and promotion of herbal drugs. Pharmacognosy Reviews, 1., 180-84.
- Siegfried ED, Jacobe G, Fatima K, (2003), recent findings on natural products with erectile-dysfunction activity. Phytochemistry, 62., 1019 – 25
- Solomon H, Man J, Wierzbicki AS, O'Brien T, Jackson G (2003), Erectile dysfunction: cardiovascular risk and the role of the cardiologist. Int J Clin Pract, 57., 96–9.
- Subramoniam, V. Madhavachandran, S. Rajasekharan, P. Pushpangadan, (1997), Aphrodisiac property of Trichopus zeylanicus extract in male mice, Journal of Ethnopharmacology, 57., 21–27

- Tajuddin, S A, Abdul L, and Iqbal A Q (2003), Aphrodisiac activity of 50% ethanolic extracts of Myristica fragrans Houtt. (nutmeg) and Syzygium aromaticum (L) Merr. &

Perry. (clove) in male mice: a comparative study, BMC Complement Altern Med, 3., 3-6

- Viqar U A, bbasi M, Hussain H, Akhtar MN, Naheed F, Choudhary MI, (2003), Phenolic glycosides from Symplocos racemosa: natural inhibitors of phosphodiesterase I, Phytochemistry, 63., 217–220.
- Ratnasooriya WD,. Fernando T.S.P, (2008), Effect of black tea brew of Camellia sinensis on sexual competence of male rats, Journal of Ethnopharmacology, 118., 373–377.
- Yakubu MT, Akanji MA, Oladiji AT. (2007), Male sexual dysfunction and methods used in assessing medicinal plants with aphrodisiac potentials, Pharmacog Rev, 1., 49-52.
- Zamble F, Sevser S , Thierry H , Bart S, Regis B, Patrick D, Claude B, Francois B, (2008), Vasoactivity, antioxidant and aphrodisiac properties of Caesalpinia benthamiana roots Alexis, Journal of Ethnopharmacology ,116., 112–119.
- Zamble, S. Sahpaza, C. Brunetb, F. Bailleula, (2008), Effects of Microdesmis keayana roots on sexual behavior of male rats, Phytomedicine, 15., 625–629.